arXiv:1208.2499v1 [astro-ph.HE] 13 Aug 2012

Phase transitions of Neutron stars and its connection with high energetic bursts in astrophysics

Ritam Mallick[a] & P K Sahu[b]

Institute of Physics, Sachivalaya Marg, Bhubaneswar 751005, INDIA

(Dated: August 14, 2012)

Abstract

Neutron stars (NS) can be made of normal hadronic matter (generally called NS) or they can have exotic states of matter in their interiors, like quark matter or color superconducting matter. Stars can have such exotic states up to their surface (called strange stars (SS)) or they can only have a quark core surrounded by hadronic matter, known as hybrid stars (HS). The hybrid stars is likely to have a mixed phase, such as both hadron and quark phases, in the intermediate region. Observational results also suggest huge surface magnetic field in certain compact stars called magnetars. The conversion of NS to SS/HS is a highly energetic process, in such conversion, the equations of state (EOS) of matter changes, resulting in the change in their masses. Special theory of relativity relates mass to energy, therefore such conversion leads to huge energy release, of the order of 10^{53} ergs. In this work we study the energy released by such conversion process. We also study the energy released by normal pulsars and magnetars. The final state of the star after conversion may be a SS or a HS, we have calculated the energy released for these final states. The energy released in the conversion of NS to SS is greater than the energy released in the conversion of NS to HS. We study the effect of magnetic field in the EOS. We find that if the magnetic field strength is of the order of 10^{14}G or more, the effect in the EOS is significant and thereby the effect of the conversion process. The energy released by magnetars is found to be more than that of normal pulsars, for NS to SS conversion process, whereas for the conversion of NS to HS the energy released by magnetars is less than normal pulsars. The amount of energy released by such conversion may only be compared to the energy observed in the gamma ray bursts (GRB).

PACS numbers: 26.60.Kp, 52.35.Tc, 97.10.Cv

---

[a] ritam.mallick5@gmail.com

[b] pradip@iopb.res.in

INTRODUCTION

The observed millisecond pulsar PSR J1614-2230 [1] recently, was found to have a very high mass, $M = 1.97 \pm 0.04$ $\mathrm{M}_{\odot}$. This value, together with the other pulsar J1903+0327 of mass $M = 1.667 \pm 0.021$ $\mathrm{M}_{\odot}$ [2] are much larger than any previous measured pulsar masses. Pulsars with such high masses, for the first time, has imposed strict constraints on determining the equations of state (EOS) of matter describing compact stars. The EOS of quark matter usually has strangeness [3–5] in them and therefore, provides an additional degree of freedom. This extra degree of freedom softens the EOS, that is, reduction of pressure for a given energy density. As a result, quark stars (QS) and hybrid stars (HS) cannot reach high value of maximum masses. However, new calculations has found that the effects from the strong interaction, such as one-gluon exchange or color-superconductivity can make the EOS stiffer and increase the maximum masses of QS and HS [6–11]. Ozel [12] and Lattimer [13] were the first to study the implications of the new mass limits from PSR J1614-2230 for QS and HS within the MIT bag model.

It has been well known for a long [14, 15] that the conversion of a neutron star (NS) to a strange star (QS) liberates huge amount of energy, simply because the star mass changes during the conversion process. From Einstein theory of relativity, mass being equivalent to energy, the mass difference is manifested in the form of energy released. The first calculation of the energy released by such conversion was proposed by Alcock et al. [14] and Olinto [15]. More detailed similar calculation for the conversion of NS to HS was done by Cheng & Dai [16] and Ma & Xie [17]. Almost all of them found the energy estimated to be greater than 10^{51}ergs, and connected them with the observed gamma ray bursts (GRB). But recent observation [18, 19] shows that the GRB to occur at cosmological distances, and so such energies are too low to power GRB at such huge distances. A systematic calculation was then done by Bombaci & Datta [20], using different models of EOS and they estimated the energy released during conversion, which is the order of 10^{53}ergs, that is in agreement with the observed GRB.

If the beaming and the γ-ray production of the GRB are quite high, then the central engine which powers such conversion process can release energy of the order of 10^{53}erg. Thereafter, various models for GRB have been proposed. Mergers of two NS or a NS and a black hole in a binary [21] being some popular models. But the recent calculation of Janka &

Ruffert [22] had shown that the energy released in such neutrino-antineutrino annihilation is much smaller to account for the GRB, thought to be occurring at large cosmological distances. So, there is still no definite answer for the central engine which powers such huge energies in the GRB.

The only other astrophysical events which can come near the energy budget of GRB are the recently observed giant flares. Three giant flares, SGR 0526-66, SGR 1900+14 and SGR 1806-20 [23] have been detected so far. The huge amount of energy in these flares can be explained by the presence of a strong magnetic field, whose strength is estimated to be larger than 10^{14}G, and such stars are named magnetars. On the other hand, theoretical calculation has suggested that such strong magnetic field may effects the gross structure of NS. A strong magnetic field affects the structure of a NS through the metric describing the star [24, 25] or by modifying the EOS of the matter through the Landau quantization of charged particles [26–30].

Therefore in this work, we plan to carry out the energy released during the conversion of NS to SS and NS to HS. The HS is likely to have a mixed phase region, and we construct a mixed phase of HS using Glendenning construction. We also aim to study the energy released for the above mentioned conversion with stars having huge magnetic field (magnetars). In the next section (Section II), we give the outline of the calculation involved in the energy released. In section III, we discuss about the EOSs which we have used. Then we discuss the magnetic field effects on the EOS in section IV. Section V is dedicated to the calculations and results. In the last section VI, we summarize our results and have a concluding discussion.

ENERGY RELEASED DURING CONVERSION

The total energy released in the conversion of NS to quark star (QS), can be either SS or HS, is actually the difference between the total binding energy of the strange star BE(QS) and the total binding energy of the neutron star BE(NS)

$$E_T = BE(QS) - BE(NS). \tag{1}$$

In terms of the gravitational mass, it can be written as difference between the gravitational mass of the NS and QS

$$E_T = [M_G(NS) - M_G(QS)]c^2, \tag{2}$$

where M_G is the gravitational mass of a star.

The total conversion energy can then be written as the sum of gravitational and internal energy change

$$E_T = E_I + E_G. \tag{3}$$

These internal and gravitational energy of conversion can be written in terms of the gravitational and internal binding energies of the NS and QS, which can further be written in terms of their respective gravitational and proper mass (or rest mass)

$$\begin{aligned} E_I &= BE_I(QS) - BE_I(NS) = [M_P(NS) - M_P(QS)]c^2, \\ E_G &= BE_G(QS) - BE_G(NS) = [M_P(QS) - M_G(QS) - M_P(NS) + M_G(NS)]c^2, \end{aligned} \tag{4}$$

where M_P is the proper mass or rest mass of a star. The proper mass and the gravitational mass of a star can be obtained by solving the structural equations for non-rotating compact stars by Tolman-Oppenheimer-Volkoff equations [31]. The gravitational and proper mass is given by

$$\begin{aligned} M_G &= \int_0^R 4\pi r^2 \epsilon(r) dr, \\ M_P &= \int_0^R dr 4\pi r^2 \Big[1 - \frac{2Gm(r)}{c^2 r}\Big]^{-1/2} \epsilon(r), \end{aligned} \tag{5}$$

where $\epsilon(r)$ being the total mass-energy density and $m(r)$ the gravitational mass enclosed within a spherical volume of radius r.

HADRONIC AND QUARK EOS

We use the non-linear relativistic mean field (RMF) model with hyperons (TM1 parameters) to describe the hadronic EOS. In this model, the baryons interact with mean meson fields [32–36].

The Lagrangian density having nucleons, baryon octet ($\Lambda, \Sigma^{0,\pm}, \Xi^{0,-}$) and leptons can be written in the form [37, 38]

$$\begin{aligned} \mathcal{L}_H = \sum_b \bar{\psi}_b [\gamma_\mu (i\partial^\mu - g_{\omega b}\omega^\mu - \frac{1}{2} g_{\rho b} \vec{\tau}.\vec{\rho}^\mu) \\ - (m_b - g_{\sigma b}\sigma)]\psi_b + \frac{1}{2}(\partial_\mu \sigma \partial^\mu \sigma - m_\sigma^2 \sigma^2) \\ - \frac{1}{4}\omega_{\mu\nu}\omega^{\mu\nu} + \frac{1}{2} m_\omega^2 \omega_\mu \omega^\mu - \frac{1}{4}\vec{\rho}_{\mu\nu}.\vec{\rho}^{\mu\nu} \end{aligned}$$

$$+\frac{1}{2}m_\rho^2\vec{\rho}_\mu.\vec{\rho}^\mu - \frac{1}{3}bm_n(g_\sigma\sigma)^3 - \frac{1}{4}c(g_\sigma\sigma)^4 + \frac{1}{4}d(\omega_\mu\omega^\mu)^2$$

$$+\sum_L \bar{\psi}_L[i\gamma_\mu\partial^\mu - m_L]\psi_L. \tag{6}$$

The leptons L are assumed to be non-interacting, whereas the baryons are coupled to the meson (scalar σ, isoscalar-vector ω_μ and isovector-vector ρ_μ). The constant parameters of the model are determined by fitting the nuclear matter saturation properties. The TM1 model fails to explain the experimentally observed strong $\Lambda\Lambda$ attraction. Mishustin & Schaffner [36] corrected the model by the adding two mesons, the isoscalar-scalar σ^* and the isovector-vector ϕ, which couple only with the hyperons. In our calculation we have used this model and denote it as TM1L.

Maintaining charge neutrality and beta equilibrium, the energy density and pressure are given by [38]

$$\varepsilon = \frac{1}{2}m_\omega^2\omega_0^2 + \frac{1}{2}m_\rho^2\rho_0^2 + \frac{1}{2}m_\sigma^2\sigma^2 + \frac{1}{2}m_{\sigma^*}^2\sigma^{*2} + \frac{1}{2}m_\phi^2\phi_0^2 + \frac{3}{4}d\omega_0^4 + U(\sigma)$$

$$+\sum_b \varepsilon_b + \sum_l \varepsilon_l\,, \tag{7}$$

$$P = \sum_i \mu_i n_i - \varepsilon \tag{8}$$

We employ the simple MIT bag model [39] to describe the quark EOS. The up, down and strange quark masses are taken to be 5MeV, 10MeV and 150MeV respectively. For the bag model, the energy density and pressure can be written as [38]

$$\epsilon^Q = \sum_{i=u,d,s} \frac{g_i}{2\pi^2}\int_0^{p_F^i} dp p^2\sqrt{m_i^2+p^2} + B_G\,, \tag{9}$$

$$P^Q = \sum_{i=u,d,s} \frac{g_i}{6\pi^2}\int_0^{p_F^i} dp \frac{p^4}{\sqrt{m_i^2+p^2}} - B_G\,, \tag{10}$$

where the Fermi momentum ${p_F}^i$ as ${p_F}^i = \sqrt{\mu_i^2 - m_i^2}$. g_i is the degeneracy of quarks i. B_G is the bag constant and is considered as a free parameter in the model.

The matter is assumed to be neutrino-free ($\mu_{\nu_e} = \mu_{\overline{\nu}_e} = 0$), and like the hadronic matter the quark matter is also charge neutral and beta equilibrium.

INCLUSION OF MAGNETIC FIELD IN EOS

The magnetic field is assumed to be along the z axis, and can be written as $\vec{B} = B\hat{k}$ [37, 40, 41], and so the charged particles get Landau quantized in the perpendicular direction to the field. Therefore the energy of the nth Landau level (n being the quantum number) [26–30, 42] takes the form

$$E_i = \sqrt{{p_i}^2 + {m_i}^2 + |q_i|B(2n + s + 1)}\,. \tag{11}$$

In this equation s is the spin of the particle and is equal to $s = \pm 1$ for the up(+) and down(-), respectively. p_i is the momentum along the field direction of particle i. Writing $2n + s + 1 = 2\nu$, the energy of the particle can be rewritten as

$$E_i = \sqrt{{p_i}^2 + {m_i}^2 + 2\nu|q_i|B} = \sqrt{{p_i}^2 + \widetilde{m}_{i,\nu}^2}\,. \tag{12}$$

Therefore, the total energy density of the hadronic matter now takes the form

$$\begin{aligned}\varepsilon = {} & \frac{1}{2}m_\omega^2\omega_0^2 + \frac{1}{2}m_\rho^2\rho_0^2 + \frac{1}{2}m_\sigma^2\sigma^2 + \frac{1}{2}m_{\sigma^*}^2\sigma^{*2} + \frac{1}{2}m_\phi^2\phi_0^2 + \frac{3}{4}d\omega_0^4 + U(\sigma) \\ & + \sum_b \varepsilon_b + \sum_l \varepsilon_l + \frac{B^2}{8\pi^2}\,,\end{aligned} \tag{13}$$

where the last term being the magnetic field contribution. The pressure can simply be represented as

$$P = \sum_i \mu_i n_i - \varepsilon\,. \tag{14}$$

For the quark matter, the thermodynamic potential in presence of strong magnetic field is written as [37, 43]

$$\Omega_i = -\frac{2g_i|q_i|B}{4\pi^2}\sum_\nu \int_{\sqrt{m_i^2+2\nu|q_i|B}}^{\mu} dE_i\sqrt{E_i^2 - m_i^2 - 2\nu|q_i|B}. \tag{15}$$

The total energy density and pressure of the strange quark matter are given by

$$\begin{aligned}\varepsilon &= \sum_i \Omega_i + B_G + \sum_i n_i\mu_i \\ p &= -\sum_i \Omega_i - B_G.\end{aligned} \tag{16}$$

EOS FOR THE MIXED PHASE

We use Glendenning conjecture [44] to determine the mixed phase. The conjecture determines the range of baryon density, where both the hadron and quark phases coexist. The stellar matter is electrically charge neutral, therefore the hadron and quark phases may be separately charged but maintaining charge neutrality in the mixed phase. The Gibbs condition determines the mechanical and chemical equilibrium between two phases, and is given by [37, 38]

$$P_{\mathrm{HP}}(\mu_e, \mu_n) = P_{\mathrm{QP}}(\mu_e, \mu_n) = P_{\mathrm{MP}}. \tag{17}$$

The volume fraction occupied by quark matter in the mixed phase is χ, and the hadronic matter is $(1-\chi)$ and is connected as

$$\chi \rho_c^{\mathrm{QP}} + (1-\chi)\rho_c^{\mathrm{HP}} = 0, \tag{18}$$

where ρ_c is respective charge density.

The energy density of the mixed phase ϵ_{MP} and the baryon density of the mixed phase n_{MP} can be calculated in terms of hadronic and quark energy density and baryon density. It is given by

$$\epsilon_{\mathrm{MP}} = \chi \epsilon_{\mathrm{QP}} + (1-\chi)\epsilon_{\mathrm{HP}}, \tag{19}$$

$$n_{\mathrm{MP}} = \chi n_{\mathrm{QP}} + (1-\chi) n_{\mathrm{HP}}. \tag{20}$$

The mixed phase lies in between the hadronic and quark phase, where both the phase coexist maintaining global charge neutrality.

RESULTS

To begin with, we would assume that a NS, with a sudden density change at the core, undergoes a phase transition. The resultant star may be a SS or a HS. We would also assume that the initial and the final star central density are same (baryonic mass is not conserved). The difference in the baryonic mass may be caused by some ejection of matter in the form of charged particles through the magnetic lines of force, which may produce jets. The conversion process may continue up to the surface or may die down after some distance. This depends on the initial energy difference between the matter phases at the center of the star and also on the initial density and spin fluctuation of the star [44].

With this assumption we then construct our initial and final star. For the initial NS, we have considered relativistic mean field EOS model for the nuclear matter. The final star may be a SS or a HS. With the above mentioned nuclear and quark EOS, we construct the mixed phase HS based on Glendenning approach. The hadronic EOS in our calculation is



FIG. 1. Pressure as a function of energy density for different EOS. The hadronic EOS is denoted by TM1L (the softer) and NL (stiffer), the quark EOS is denoted as quark, and the mixed phase EOS is denoted as mix.

$NS \rightarrow SS$	n_b/n_0	$M_G(NS)$	$M_P(NS)$	$M_G(SS)$	$M_P(SS)$	E_G	E_I	E_T
$TM1L \rightarrow bag145$	6	1.695	1.868	1.845	2.154	2.43	-5.11	-2.68
	8	1.648	1.809	1.897	2.231	3.10	-7.55	-4.44
$TM1L \rightarrow bag170$	6	1.695	1.868	1.22	1.32	-1.3	9.8	8.5
	8	1.648	1.809	1.37	1.501	-0.54	5.51	4.97
$NL \rightarrow bag145$	6	2.637	3.177	1.845	2.154	-4.14	18.3	14.16
	8	2.63	3.16	1.897	2.231	-3.5	16.61	13.11
$NL \rightarrow bag170$	6	2.637	3.177	1.22	1.32	-7.87	33.21	25.34
	8	2.63	3.16	1.37	1.501	-7.13	29.67	22.54
$NS \rightarrow HS$								
$TM1L \rightarrow mix170$	6	1.695	1.868	1.23	1.33	-1.31	9.62	8.31
	8	1.648	1.809	1.38	1.51	-0.56	5.35	4.79
$NL \rightarrow mix170$	6	2.637	3.177	1.23	1.33	-7.87	33.03	25.16
	8	2.63	3.16	1.38	1.51	-7.15	29.51	22.36

TABLE I. Table showing the energy released during the phase transition from NS to SS/HS for the non magnetic star. The masses are in terms of solar masses and the energies in terms of 10^{53}ergs.

fixed, but we can regulate the quark EOS by changing the bag constant. The masses of the light quarks are quite bounded and we take them to be 5MeV (u) and 10MeV (d). The mass of s-quark is still not well established, but expected to lie between $100-300$MeV. Therefore, we have kept the mass of the s-quark fixed at 150MeV. We take different bag constant (B_G) to have different quark matter EOSs. For simplicity, we will denote ${B_G}^{1/4} = 170 MeV = B_g$. For the mixed phase EOS, a mixed phase region with constant bag pressure less than 170 MeV is not possible, having mass of s-quark 150MeV. Thus, we have shown results with bag pressure 170MeV for SS and HS. With such bag pressure it is impossible to produce massive compact star, therefore, for comparison we have also shown results for SS with bag pressure 145MeV. For comparison, we have also shown results with much stiffer hadronic EOS, non linear walecka (NL) [45]. In figure 1, we have plotted pressure as a function of energy density for different EOSs. The hadronic EOS is moderately stiff and the MIT EOS with bag 145MeV is quite stiff, whereas the MIT EOS with bag pressure 170MeV is quite soft. The mixed phase EOS with bag pressure 170MeV goes from the hadronic to quark EOS.

Table I shows the energy released by the conversion of NS to SS. For the first set of EOSs, with RMF and bag 145MeV, the energy released is negative. This is because the final star is more massive than the initial star, therefore, such conversion is not possible. This conversion process is energetically restricted. For bag pressure 170MeV, we find that such conversion of NS to SS is possible, and the energy released is few times 10^{53}ergs. This conversion process is energetically possible because the initial star is more massive than final star and the energy released during the conversion of hadronic matter to quark matter drives the conversion process. We also find that with the increase in central density from six times normal nuclear density ($6n_0$) to eight times normal nuclear density ($8n_0$), there is a fall in energy released. This is because, the hadronic EOS reaches maximum mass at much less density than the quark EOS. As mentioned earlier, we have used a stiffer hadronic EOS (NL) for comparison. With such hadronic EOS, the conversion of NS to SS with bag 145MeV is possible, as the energy released in this case is positive. This is because the initial hadronic star has greater mass than the final star. The mass difference is positive, which drives the conversion process. The energy released for this case is much higher and is the order of 10^{54}ergs. As we go to higher central density, the energy released decreases because at such densities the hadronic stars reaches maximum mass but for the quark EOS the maximum



FIG. 2. Pressure against energy density for different EOS without and with magnetic field, with field strength 2×10^{14}G.

$NS \rightarrow SS$	n_b/n_0	$M_G(NS)$	$M_P(NS)$	$M_G(SS)$	$M_P(SS)$	E_G	E_I	E_T
$TM1L \rightarrow bag145$	6	1.684	1.855	1.747	2.026	1.94	-3.06	-1.12
	8	1.638	1.796	1.803	2.104	2.56	-5.51	-2.95
$TM1L \rightarrow bag170$	6	1.684	1.855	0.93	1.04	-1.1	14.58	13.48
	8	1.638	1.795	1.15	1.314	0.13	8.6	8.73
$NS \rightarrow HS$								
$TM1L \rightarrow bag170$	6	1.684	1.855	1.28	1.38	-1.27	8.49	7.22
	8	1.638	1.796	1.385	1.515	0.5	5.02	4.52

TABLE II. Table showing the energy released during the phase transition from NS to SS/HS with magnetic field of 2×10^{14}G. The masses are in terms of solar masses and the energies in terms of 10^{53}ergs.

mass is still not reached. It is the characteristic of the hadronic EOS that it saturates at lower densities. Next we calculate the conversion of NS to HS. As stated earlier, it is not possible to construct mixed phase HS with bag 145MeV, we have shown results only for bag 170MeV. The qualitative result remains the same, and the energy released is slightly lower. For simple TM1L NS to HS, the energy released is of the order of 10^{53}ergs and for NL hadronic EOS the energy released is 10^{54}ergs.

Next we consider magnetic field in the EOS. The effect of magnetic field is insignificant if the field strength is less than 10^{14}G. Magnetic field makes the curve softer due to the negative effect of Landau quantization on the matter pressure and the positive effect on the matter energy density. Such magnetic field have almost no effect on the hadronic EOS, but has significant effect on the quark EOS. The EOS for mixed HS in the low density region is not much affected, however in the high density region its change is significant. The magnetic field makes the mixed phase EOS flat, that is in the intermediate region the pressure is larger than the non magnetic EOS, that is in the intermediate region the EOS is stiff but it become soft later in the higher densities. In fig 2, we have shown EOS curves with and without magnetic field, for the quark and mixed phase EOS. The magnetic field therefore makes the EOS for the quark matter soft, which thereby makes the mass-radius curve flat. Therefore, for a fixed central density the mass of the SS having magnetic field is much less than that of the SS without magnetic field. The surface field strength observed in magnetars lie between 10^{14}G to 10^{15}G. Therefore in our calculation, we have taken the effect of magnetic field in the EOS, with field strength 2×10^{14}G. With the magnetic field and bag pressure being constant throughout the star, higher surface magnetic field makes the mixed phase region to disappear. As shown in Table II, for the first set of EOSs, with simple TM1L hadronic model and bag model with bag constant 145MeV in the quark matter, the energy released is negative and the conversion process is energetically restricted, we therefore would move to the set with bag constant 170MeV in the quark matter. The energy released lies between $10^{53} - 10^{54}$ergs. Comparing with the non magnetic case, the energy released is higher ($\sim 60 - 70\%$). This is because, for a fixed central density, the mass of the SS is much lower than that of the non magnetic star. The mass of the NS does not change much with the magnetic field. For the NS to HS conversion with the magnetic field the energy released is lesser than the NS to SS ($\sim 50\%$). This is because for the mixed phase EOS, in the intermediate density range the EOS is stiffer than the non magnetic counterpart, making the mass of the magnetic HS higher than the non magnetic HS. The energy released in the conversion process of NS to HS having magnetic field effect does not change much in comparison to the non magnetic conversion process ($\sim 5 - 10\%$). The magnetic field have slight effect on mixed phase EOS. Such low energy value can also be responsible for the fact that the conversion process cannot sustain up to the star surface, and stops inside the star, resulting in the HS having both quark region (conversion possible) and hadronic region (conversion stops).

The previous calculation done by bombaci & datta [20], assumed that the baryonic mass of the star to remains unaltered. In their work the energy released for such a conversion process is always about few times 10^{53}ergs. With the same baryonic mass, the NS has always greater mass than the final SS or HS. Therefore energetically such conversion process are always possible.

We therefore, find that there is conceptually some basic difference between the two assumptions. Although for most of the cases, with the two assumptions the energy released is around 10^{53}ergs, the qualitative results are different. For the fixed baryonic mass assumptions the energy released is always positive and is always few times 10^{53}ergs. For the fixed central energy density the conversion process is always not possible and it is energetically restricted if the final mass of the star is greater than the initial mass, that is, if the quark EOS is much stiffer than the hadronic EOS. Also for the fixed central density assumption, the conversion process is accompanied by particle or matter ejection. The main result from our calculation is that the energy released in the conversion of NS to SS/HS is in the range $10^{53} - 10^{54}$ergs. For some cases the conversion process is energetically restricted. The energy released in the conversion of NS to SS is always greater than the energy released during conversion of NS to HS. This huge amount of energy release is only comparable to the energy observed during the GRB. As the detailed picture of the conversion mechanism is not known, we have no idea of its true nature. But we may guess that magnetars would release the energy more efficiently from their magnetic poles than the normal NS. We also are unsure about the nature of the magnetic field after the conversion process. The main result of our calculation is that the energy released in such conversion processes depends on the initial and final star masses, and therefore a initial NS cannot result to final more massive SS or HS.

SUMMARY AND CONCLUSION

In this work, we have calculated the energy released during the conversion of NS to SS or HS. The total energy released is the sum of the gravitational and internal energy of conversion. Our first assumption was that the conversion process starts with a sudden spin down of the star, resulting in a huge density fluctuation at the core, initiating the phase transition. The baryonic mass is not conserved, and the central density remains same after the conversion process. The difference in the baryonic mass can be restored by assuming some mechanism for mass ejection of charged particles through the magnetic lines of force, producing jets. The conversion process may continue up to the surface or may die out after some distance. This depends on the energy difference between the matter phases at the center of the star and also on the initial density and spin fluctuation of the star. For the NS we have considered relativistic mean field EOS model for the hadronic matter. For the quark matter (SS or a HS), we have considered simple MIT bag model. For the HS we have considered mixed phase region, between the outer hadron and inner quark matter. We have used two bag pressures in our calculation, 145 and 170MeV. For comparison, we have also used the stiff NL hadronic EOS. As it is not possible to construct mixed phase HS with the given set of EOS having bag constant 145MeV, therefore, for HS we have shown results only for bag constant 170MeV.

We find that if the mass of the final star is greater than the initial NS, the conversion process is energetically restricted. Therefore, a initial NS cannot produce more massive SS/HS. We also find that for most of the cases the energy released is of the order of 10^{53}ergs. For the stiff hadronic EOS the energy released is much higher. The energy released for the conversion of NS to HS is almost same as that of the energy released in the conversion of NS to SS.

We have also studied the conversion of NS to SS/HS both having high surface magnetic field (observed magnetars). The effect of magnetic field is insignificant if the field strength is less than 10^{14}G. We have shown result with magnetic field 2×10^{14}G, as for a fixed magnetic field, we do not get a mixed phase region with higher magnetic field. Magnetic field makes the curve less stiffer due to the negative effect of Landau quantization on the matter pressure and the positive effect on the matter energy density. The effect of the magnetic field on the EOS for the hadronic matter is not very much but for quark matter it is quite significant. Therefore, the change in mass of the hadronic star and HS is not much, whereas the change in the mass of SS is of considerable amount. The energy released during the conversion of NS to SS is slightly higher, than the non magnetic field induced EOS. The energy released in the conversion of NS to HS is less than the conversion of NS to SS, which can also be responsible for the fact that the conversion process cannot sustain up to the star surface, and stops inside the star, resulting in a HS having both quark region (conversion happens) and hadronic region (conversion stops).

Assuming the baryonic mass of the star to remain unaltered, Bombaci and Dutta [20] performed their calculation and found that the NS has always greater mass than the final SS. Therefore energetically, such conversion process are always possible. There is significant difference between the two assumptions. The energy released is quite different. With the assumption of fixed baryon mass, the energy released is always few times 10^{53}ergs. But with assumption of fixed central density the energy released can be as high as 10^{54}ergs. Also for the fixed central density assumption the conversion process is accompanied by particle or matter ejection. The magnetic field makes the quark EOS much softer, and so the energy released in the conversion of NS to SS/HS is larger than the non magnetic field. As the actual process takes place at cosmological distances, we do not have a better understanding of such phenomena. We are not sure about what fraction of the energy released is used in initiating and maintaining the conversion process. As there are no other source of external energy available to the star, other than the rotational energy, a huge amount of energy may be used in the actual conversion process. We can only conclude that if a small fraction of the energy of the conversion process is released it may manifest itself at least in the form of giant flares, that are associated with the magnetars. If a substantial amount of energy is released, it may answer for the energies of the GRB. Therefore, a detailed study, theoretical and observational, is still needed for the pulsars and magnetars to have a understanding of the energetic of the conversion process.

[1] Demorest, P., Pennucci, T., Ransom, S., Roberts, M., & Hessels, J. 2010, Nature **467**, 1081

[2] Freire, P. C. C., Bassa, C., Wex, N., et al. 2010, MNRAS, **412**, 2763

[3] Itoh, N. 1970, Prog. Theor. Phys. **44**, 291

[4] Bodmer, A. R. 1971, Phys. Rev. D **4**, 1601

[5] Witten, E. 1984, Phys. Rev. D **30**, 272

[6] Rüster, S. B. & Rischke, D. H. 2004, Phys. Rev. D **69**, 045011

[7] Horvath, J. E. & Lugones, G. 2004, Astron. & Astrophys. **442**, L1

[8] Alford, M., Blaschke, D., Drago, A., et al. 2007, Nature, **445**, 7

[9] Fischer, T., Sagert, I., Pagliara, G., et al. 2010, AstroPhys. J Supp. **194**, 39

[10] Kurkela, A., Romatschke, P., & Vuorinen, A. 2010, Phys. Rev. D **81**, 105021

[11] Kurkela, A., Romatschke, P., Vuorinen, A., & Wu, B. 2010, arXiv:1006.4062

[12] Özel, F., Psaltis, D., Ransom, S., Demorest, P., & Alford, M. 2010, AstroPhys. J Lett. **724**, L199

[13] Lattimer, J. M. & Prakash, M. 2010, arXiv:1012.3208

[14] Alcock, C., Farhi, E. and Olinto, A. 1986, AstroPhys. J **310**, 261

[15] Olinto, A. 1987, Phys. Lett. B **192**, 71

[16] Cheng, K. S. & Dai, Z. G. 1996, Phys. Rev. Lett. **77**, 1210

[17] Ma, F. & Xie, B. 1996, Astrophys. J **527**, L51

[18] Kulkarni, S. R., et al. 1998, Nature **393**, 35

[19] Kulkarni, S. R., et al. 1999, Nature **398**, 389

[20] Bombaci, I. & Datta, B. 2000, Astrophys. J **530**, L69

[21] Paczyñski, B. 1998, Astrophys. J **494**, L45

[22] Janka, H. -Th. & Rufert, M. 1996, Astron. & Astrophys. **307**, L33

[23] Palmer, D. M., Barthelmy, S., Gehrels, N. et al. 2005, Nature bf 434, 1107

[24] Bocquet, M., Bonazzola, S., Gourgoulhon, E., Novak, J. 1995, Astron. & Astrophys. **301**, 757

[25] Cardall, C. Y., Prakash, M. & Lattimer, J. L. 2001, AstroPhys. J **554**, 322

[26] Chakrabarty, S., Bandyopadhyay, D. & Pal, S. 1997, Phys. Rev. Lett. **78**, 2898

[27] Yuan, Y. F. & Zhang, J. L. 1999, AstroPhys. J **25**, 950

[28] Broderick, A., Prakash, M. & Lattimer, J. M. 2000, AstroPhys. J **537**, 351

[29] Chen, W., Zhang, P. Q. & Liu, L. G. 2005, Mod. Phys. Lett. A **22**, 623

[30] Wei, F. X., Mao, G. J., Ko, C. M., Kissinger, L. S., Stoecker, H. & Greiner, W. 2006, Jour. Phys. G **32**, 47

[31] Shapiro, S. L. & Teukolsky, S. A. *Black Holes, White Dwarfs, and Neutron Stars*, (John Wiley & Sons, New York, 1983)

[32] Boguta, J. & Bodmer, R. A. 1977, Nucl. Phys. A **292**, 413

[33] Glendenning, N. K. & Moszkowski, S. A. 1991, Phys. Rev. Lett. **67**, 2414

[34] Ghosh, S. K., Phatak, S. C. & Sahu, P. K. 1995, Z. Phys. A **352**, 457

[35] Sugahara, Y. & Toki, H. 1994, Nucl. Phys. A **579**, 557

[36] Schaffner, J. & Mishustin, I. N. 1996, Phys. Rev. C **53**, 1416

[37] Mallick, R. & Sahu, P. K. 2012, arXiv:1207.4780

[38] Mallick, R. 2012, arXiv:1207.4782

[39] Chodos, A., Jaffe, R. L., Johnson, K., Thorn, C. B. and Weisskopf, V. F. 1974, Phys. Rev. D **9**, 3471

[40] Sinha, M. & Mukhopadhyay, B. 2010, arXiv:1005.4995

[41] Mallick, R. & Sinha M. 2011, Mon. Not. R. Astron. Soc. **414** , 2702

[42] Landau, L. D. & Lifshitz, E. M. *Quantum Mechanics*, (Pergamon Press, Oxford, 1965)

[43] Chakrabarty, S. & Sahu, P. K. 1996, Phys. Rev. D **53**, 8

[44] Glendenning, N. K. 1992, Phys. Rev. D **46**, 1274

[45] Ellis, J., Kapusta, J. I. & Olive, K. A. 1991, Nucl. Phys. B **348**, 345